

PROFESSIONAL INFORMATION

SCHEDULING STATUS: **S0**

Category D: Complementary medicine.

Discipline: Discipline Specific

33.6 Western Herbal Medicine

PROPRIETARY NAME, STRENGTH, PHARMACEUTICAL FORM –

CITRICIDAL LIQUID® , 30 ml Liquid

COMPOSITION

Each one (1) drop of CITRICIDAL LIQUID® contains the following actives:

Citricidal® standardized extract of Grapefruit (*Citrus 25 mg
parasis*) [Seed and Pulp Extract]

Inactive Ingredients: Citric acid, glycerine, lactic acid

This product is suitable for Vegetarians and non-GMO. (Sugar Free)

PHARMACOLOGICAL CLASSIFICATION

Complementary Medicine

INDICATION & PHARMACOLOGICAL ACTION

CITRICIDAL LIQUID® contributes to the microbial balance in the body organs and tissues and supports a healthy immune system. CITRICIDAL LIQUID® also provides antioxidants that protects cells against oxidative damage caused by free radicals.

Ingredients	Uses for ingredient
Citricidal®	Contains bioactive compounds which are associated with therapeutic properties including antiallergic, antiatherogenic, anti-inflammatory, antimicrobial, anticarcinogenic, antithrombotic, cardioprotective, and vasodilatory effects

CONTRA INDICATIONS:

If you are taking any chronic medication, do not use this product without consulting your healthcare professional. Do not take CITRICIDAL LIQUID® if you are hypersensitive (allergic) to any of the ingredients. If you have allergies, be sure to check with your healthcare professional before taking this supplement.

WARNING AND SPECIAL PRECAUTIONS

Should you have any of the following diagnosed conditions, consult with your healthcare professional before use: heart muscle disorder (cardiomyopathy), Hormone sensitive cancers and conditions, Irregular heartbeat, Postmenopausal adults, A rare inherited blood condition called porphyria, Kidney disorders, Seizure disorders, including epilepsy, High bilirubin levels in the blood. If you are taking the following medication, please consult with your healthcare practitioner before taking: Artemether (Artemam, Paluther), Busiprone (BuSpar), Carbamazepine (Tegretol), Carvedilol (Coreg), Estrogen, high blood pressure (Calcium channel blockers), Quinidine, Sedative medications (Benzodiazepines), Methadone (Dolophine), Medications used for lowering cholesterol (Statins), Caffeine, Warfarin (Coumadin), Medications used to prevent seizures (Anticonvulsants), Sedative medications (CNS depressants). If you are taking the following medication, please consult with your healthcare professional before taking: digoxin, sodium polystyrene sulfonate, tetracycline/quinolone antibiotics (such as tetracycline, ciprofloxacin). If any of the following happens, stop using CITRICIDAL LIQUID® and tell your doctor or go to the casualty department at your nearest hospital:

- Allergic reactions— rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing;



These are all very serious effects. If you have them, you may have had a serious reaction to CITRICIDAL LIQUID®. You may need urgent medical attention or hospitalisation.

Side effects that usually do not require medical attention (report to your healthcare professional if they continue):

- nausea/vomiting, constipation, loss of appetite, mental/mood changes, unusual tiredness.

These are all very serious effects. If you have them, you may have had a serious reaction to CITRICIDAL LIQUID®. You may need urgent medical attention or hospitalisation.

Seek advice from a healthcare professional if you have any medical condition. Discontinue use immediately should any adverse reaction occur.

Read this leaflet carefully because it contains important information for you.

This product is available without a doctor's prescription.

- Keep this leaflet. You may need to read it again.
- Do not share medication with any other person.
- Ask your pharmacist if you need more information or advice.
- You must consult a doctor should your condition worsen or does not improve.

DO NOT USE THIS PRODUCT:

- If you are hypersensitive (allergic) to any of the ingredients listed.
- Do not accept this package if seals are broken.

INTERACTIONS

Always tell your healthcare professional if you are taking any other medicine. It is possible that exposure of certain supplement ingredients may interfere with certain medications. (See **Warnings & Special precautions**)

DOSAGE AND DIRECTIONS FOR USE

Adults: Mix 1 to 5 drops in a glass of water or juice, one to three times daily or as directed by your healthcare practitioner.

Do not take more than the recommended dose.

SIDE EFFECTS

Should your general health worsen, or if you experience any untoward effects while being exposed to this product, please consult your doctor, pharmacist or other healthcare professional for advice.

KNOWN SYMPTOMS OF OVER DOSAGE AND PARTICULARS OF ITS TREATMENT

See side-effects.

Treatment:

Discontinue use and consult your doctor, pharmacist or other healthcare professional for advice.

STORAGE INSTRUCTION

Store in a cool, dark place, below 25 °C.

Protect from direct sunlight and heat.

Store in the original bottle.

Keep the bottle tightly closed.

Replace the protective cap after each use.

Do not use after the expiry date stated on the bottle.

Do not use CITRICIDAL LIQUID® if you notice visible signs of deterioration.

IDENTIFICATION

Liquid.

PRESENTATION

A 30 ml HDPE bottle with LDPE dropper and lid.

REGISTRATION NUMBER

To be allocated by SAHPRA upon registration.

This medicine has not been evaluated by SAHPRA for quality, safety, or intended use.

NAME AND BUSINESS ADDRESS OF APPLICANT

Distributed by:

Amipro Advanced Development Products (Pty) Ltd

Unit 3, Eastgate Business Park, 1 South Road, Eastgate Extension,

Sandton, 2146

Tel: 011-802-8101

DATE OF PUBLICATION

To be allocated by the regulatory authority.

PROFESIONELE INLIGTINGSBLAD

SKEDULERINGSSTATUS: S0

Kategorie D: Komplementêre medisyne.

Dissipline: Dissipline Spesifiek

33.6 Westerse Krui Medisyne

HANDELSNAAM EN DOSEERVORM,

CITRICIDAL LIQUID[®], 30 ml Vloeistof

SAMESTELING

Elke (1) druppel CITRICIDAL LIQUID[®] bevat die volgende aktiewe stowwe:

Sitroedodende* gestandaardiseerde uittreksel van 25 mg
pomelo (*Citrus parviflora*) [Saad ekstrak]

Onaktiewe bestanddele: : Sitroensuur, glierien, melksuur

Hierdie produk is geskik vir vegetêriërs en nie-GGO. (Suikervry)

FARMAKOLOGIESE KLASIFIKASIE

Komplementêre medisyne

AANDUIDING EN FARMAKOLOGIESE WERKING

CITRICIDAL LIQUID[®] dra by tot die mikrobiële balans in die liggaamsorgane en weefsels en ondersteun 'n gesonde immuunstelsel. CITRICIDAL LIQUID[®] verskaf ook antioksidante wat selle beskerm teen oksidatiewe skade wat deur vrye radikale veroorsaak word.

Bestanddele	Gebruik vir bestanddeel
Sitroedodend*	Bevat bioaktiewe verbindings wat geassosieer word met terapeutiese eienskappe, insluitend anti-allergene, anti-aterogene, anti-inflammatoriese, antimikrobiële, antikarsinogene, antitrombotiese, kardiobeskerende en vasodilerende effekte

KONTRA-INDIKASIES:

As jy enige chroniese medisyne neem, moenie hierdie produk gebruik sonder om jou gesondheidsorgpraktisyn te raadpleeg nie. Moenie CITRICIDAL LIQUID[®] neem as jy hipersensitief (allergies) is vir enige van die bestanddele nie. As jy allergieë het, maak seker dat jy met jou gesondheidsorgpraktisyn gaan voordat jy hierdie aanvulling neem.

WAARSKUWING EN SPESIALE VOORSORGMATREËLS

Indien jy enige van die volgende gedagnoseerde toestande het, raadpleeg jou gesondheidsorgpraktisyn voor gebruik: hartspierversteuring (kardiomiopatie), Hormoonsensitiewe kankers en toestande, Onreëlmatige hartklop, Postmenopousale volwassenes, 'n Seldsame oorerflikke bloedtoestand genaamd porfirie, Nierafwykings, Beslagleggingsversteurings, insluitend epilepsie, Hoë bilirubienvlakke in die bloed.

As jy die volgende medisyne neem, raadpleeg asseblief jou gesondheidspraktisyn voordat jy neem: Artemether (Artemam, Paluther), Buspiron (BuSpar), Carbamazepine (Tegretol), Carvedilol (Coreg), Estrogeen, hoë bloeddruk (Kalsiumkanaalblokkers), Quindine, Kalmerende medikasie (Bensodiasepiene), Metaadon (Dolophine), medisyne wat gebruik word vir die verlagings van cholesterol (Statiene), Kafêien, Warfarin (Coumadin), medisyne wat gebruik word om aanvalle te voorkom (Antikonvulsante), Kalmerende medisyne (SSS-depressante). As enige van die volgende gebeur, hou op om CITRICIDAL LIQUID[®] te gebruik en vertel jou dokter of gaan na die ongevalle-afdeling by jou naaste hospitaal:

- Allergiese reaksies- uitslag, jeuk / swelling (veral van die gesig / tong / keel), erge duiseligheid, probleme met asemhaling;

Dit is alles baie ernstige gevolge. As jy dit het, het jy dalk 'n ernstige reaksie op CITRICIDAL LIQUID[®] gehad. Jy sal dalk dringende mediese hulp of hospitalisasie nodig hê.

Nuwe-effekte wat gewoonlik nie mediese hulp benodig nie (rapporteer aan jou gesondheidsorgpraktisyn as hulle voortduur):

- naarheid / braking, hardlywigheid, verlies aan eetlus, geestelike / gemoedsveranderinge, ongewone moegheid.

Dit is alles baie ernstige gevolge. As u dit het, het u moontlik 'n ernstige reaksie op CITRICIDAL LIQUID[®] gehad. Miskien het u dringende mediese hulp of hospitalisasie nodig.

Raadpleeg 'n gesondheidsorgpraktisyn as u enige mediese toestand het. Staak gebruik onmiddellik indien enige nadelige reaksie plaasvind.

Lees hierdie inligtingsblad aandagtig deur, want dit bevat belangrike inligting vir u. Hierdie produk is beskikbaar sonder doktersvoorskrif.

- Hou hierdie inligtingsblad. Miskien moet u dit weer lees.
- Moenie medisyne met enige ander persoon deel nie.
- Vra jou apteker of jy meer inligting of advies benodig.
- Jy moet 'n dokter raadpleeg indien jou toestand vererger of nie verbeter nie.

MOENIE HIERDIE PRODUK GEBRUIK NIE:

• As jy hipersensitief (allergies) is vir enige van die bestanddele wat gelys word.

• Jy moet nie hierdie produk aanvaar as seëls gebreek word nie.

INTERAKSIES

Vertel altyd jou gesondheidsorgpraktisyn as jy enige ander medisyne gebruik. Dit is moontlik dat blootstelling van sekere aanvullingsbestanddele met sekere medisyne 'n interaksie mag hê. (Sien **Waarskuwings en Spesiale voorsorgmatreëls**)

DOSIS EN AANWYSINGS VIR GEBRUIK

Volwassenes: Meng 1 tot 5 druppels in 'n glas water of sap, een tot drie keer per dag of soos voorgeskryf deur jou gesondheidspraktisyn. **Moenie meer as die aanbevole dosis neem nie.**

NEWE-EFFEKTE

Indien jou algemene gesondheid vererger of as jy enige nadelige effekte ervaar tydens die gebruik van hierdie medisyne, raadpleeg asseblief jou dokter, apteker of ander gesondheidsorgpraktisyn vir raad. As jy enige nuwe-effekte sien wat nie in hierdie inligtingsblad genoem word nie, stel asseblief jou dokter, apteker of gesondheidsorgpraktisyn in kennis.

BEKENE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING DAARVAN

Sien nuwe-effekte.

Behandeling:

Staak gebruik en raadpleeg jou dokter, apteker of ander gesondheidsorgpraktisyn vir advies.

BERGING INSTRUKSIE

Bêre in 'n koel, donker plek, onder 25 °C.

Beskerms teen direkte sonlig en hitte.

Bêre n die oorspronklike bottel.

Hou die bottel styf toe.

Vervang die beskermerende dop na elke gebruik.

Moet nie gebruik word na die vervaldatum wat op die bottel aangedui word nie.

Moenie CITRICIDAL LIQUID[®] gebruik as u sigbare tekens van agteruitgang opmerk nie.

IDENTIFIKASIE

Vloeistof.

AANBIEDING

n 30 ml HDPE-bottel met LDPE-druppers en deksel.

REGISTRASIE NOMMER

Om by registrasie deur SAHPRA toegeken te word.

Hierdie medisyne is nie deur SAHPRA geëvalueer vir kwaliteit, veiligheid of beoogde gebruik nie.

NAAM EN BESIGHEIDSADRES VAN APPLICANT

Versprei deur: Amipro Advanced Development Products (Pty) Ltd
Unit 3, Eastgate Business Park, 1 South Road,
Eastgate Extension,
Sandton, 2146,
Tel: 011- 802-8101

DATUM VAN PUBLIKASIE

Om deur die regulerende owerheid toegeken te word.