

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S0

Category D: Complementary medicine.

Discipline: Discipline Specific

33.4 Traditional Chinese Medicine

PROPRIETARY NAME, STRENGTH, PHARMACEUTICAL FORM –

SERENAGEN® , 60 & 120 Tablets

READ ALL OF THIS LEAFLET CAREFULLY BECAUSE IT CONTAINS

IMPORTANT INFORMATION FOR YOU

- **SERENAGEN®** is available without a doctor's prescription.
- Nevertheless, you still need to use **SERENAGEN®** carefully to get the best results from it.
- Keep this leaflet. You may need to read it again.
- Do not share **SERENAGEN®** with any other person.
- Ask your pharmacist or healthcare professional if you need more information or advice.

WHAT IS IN THIS LEAFLET

- What **SERENAGEN®** is and what it is used for.
- What you need to know before you take **SERENAGEN®**.
- How to take **SERENAGEN®**.
- Possible side effects.
- How to store **SERENAGEN®**.
- Contents of the pack and other information.

WHAT SERENAGEN® CONTAINS

Each two (2) tablets of **SERENAGEN®** contains the following actives:

Rehmannia (Rehmannia glutinosa) [Root 4:1 extract]	290 mg
Schisandra (Schisandra chinensis) [Fruit 4:1 extract]	120 mg
Jujube (Ziziphus spinosa) [Fruit 4:1 extract]	120 mg
Chinese Asparagus (Asparagus cochinchinensis) [Root 4:1 extract]	70 mg
Dong Quai (Angelica sinensis) [Root 4:1 extract]	70 mg
Ophiopogon (Ophiopogon japonicus) [Root 4:1 extract]	70 mg
Scrophularia (Scrophularia ningpoensis) [Root 4:1 extract]	70 mg
Asian Ginseng (Panax ginseng) [Root 4:1 extract]	40 mg
Chinese Salvia (Salvia miltiorrhiza) [Root 4:1 extract]	40 mg
Poria Fungus (Wolfiporia cocos) [Whole 4:1 extract]	40 mg
Polygala (Polygala tenuifolia) [Root 4:1 extract]	40 mg
Platycodon (Platycodon grandiflorum) [Root 4:1 extract]	30 mg

Inactive Ingredients: Microcrystalline cellulose, croscarmellose sodium, cellulose, stearic acid (vegetable), silica, magnesium stearate (vegetable), and coating (hypromellose, medium-chain triglycerides, and hydroxypropylcellulose).

This product is suitable for Vegetarians, is non-GMO, gluten and dairy free. (Sugar Free)

WHAT SERENAGEN® IS AND WHAT IT IS USED FOR

SERENAGEN® offers a unique combination of adaptogenic herbs traditionally used to support organs and systems involved in the stress response. Formulated for individuals who are stressed and tense.

BEFORE YOU TAKE SERENAGEN®

If you are taking any chronic medication, do not use this product without consulting your healthcare provider. Do not take **SERENAGEN®** if you are hypersensitive (allergic) to any of the ingredients. If you have allergies, be sure to check with your healthcare professional before taking this supplement. **Surgery:** **SERENAGEN®** might affect blood sugar levels. This might interfere with blood sugar control during and after surgery. Stop using **SERENAGEN®** at least 2 weeks before a scheduled surgery. **Latex allergy:** People who are allergic to latex might also be allergic to **SERENAGEN®**.

This medicine has not been evaluated by the SAHPRA for quality, safety, or intended use.

PREGNANCY AND BREASTFEEDING

Safety during pregnancy and breastfeeding has not been established, consult with your healthcare professional before use. If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult with your healthcare professional before taking this supplement.

DRIVING AND USING MACHINES

It is not always possible to predict to what extent **SERENAGEN®** may interfere with the daily activities of a patient. Patients should ensure that they do not engage in activities relating to requiring mental alertness, judgement and/or sound co-ordination and vision e.g. driving, riding, flying, sailing, operating machines/equipment, until they are aware of the measure to which **SERENAGEN®** may affect you.

TAKING OTHER MEDICINES WITH SERENAGEN®

Complementary medicine products contain ingredients which may interact with some medication. If you are taking any chronic medication, do not use this product without consulting your medical practitioner.

HOW TO TAKE SERENAGEN®

Do not share medicines prescribed for you with any other person. Always take **SERENAGEN®** exactly as described in this leaflet, or as your doctor, healthcare provider has instructed you. You should check with your doctor, pharmacist or healthcare provider if you are unsure.

Directions for use:

Adults: Take two (2) tablets daily or as directed by your healthcare practitioner.

Do not take more than the recommended dose.

IF YOU TAKE MORE SERENAGEN® THAN YOU SHOULD

In the event of overdose, consult your doctor, pharmacist. Should none of these be available, contact the nearest hospital or poison control centre.



IF YOU FORGET TO TAKE SERENAGEN®

Do not take a double dose to make up for a missed dose.

POSSIBLE SIDE EFFECTS

SERENAGEN® may have side effects.

Should you have any of the following diagnosed conditions, consult with your healthcare professional before use: Diabetes, high blood pressure, Bleeding disorders, Hormone-sensitive conditions such as breast cancer, uterine cancer, ovarian cancer, endometriosis, or uterine fibroids. If you are taking the following medication, please consult with your healthcare practitioner before taking: diabetes (Antidiabetes drugs), high blood pressure (Antihypertensive drugs), Bleeding disorders, slow blood clotting (Anticoagulant / Antiplatelet drugs). If any of the following happens, stop using **SERENAGEN®** and tell your doctor or go to the casualty department at your nearest hospital:

- Allergic reactions— rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

These are all very serious effects. If you have them, you may have had a serious reaction to **SERENAGEN®**. You may need urgent medical attention or other healthcare professional for advice.

Side effects that usually do not require medical attention (report to your healthcare professional if they continue):

- nausea/vomiting, constipation, loss of appetite, increased thirst, increased urination, mental/mood changes, unusual tiredness.

Not all side effects and interactions reported for **SERENAGEN®** are included in this leaflet.

Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your doctor, pharmacist or other healthcare professional for advice. If you notice any side effects not mentioned in this leaflet, please inform your doctor, pharmacist or healthcare professional.

REPORTING OF SIDE EFFECTS

If you experience side effects, talk to your doctor, pharmacist or other healthcare professional. You can also report side effects to SAHPRA via "6.04 Adverse Drug Reaction Reporting Form", found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of **SERENAGEN®**.

STORAGE AND DISPOSING OF SERENAGEN®

Store all medicines out of the reach of children.

Store in a cool, dark place, below 25 °C.

Protect from direct sunlight and heat.

Store in the original packaging.

Do not use after the expiry date stated on the bottle.

Do not use **SERENAGEN®** if you notice visible signs of deterioration.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

PRESENTATION OF SERENAGEN®

60 TABLETS - A white 175 ml HDPE bottle with a white cap, induction seal, tamper shrink seal, wadding and silica gel sachet.

120 TABLETS - A white 250 ml HDPE bottle with a white cap, induction seal, tamper shrink seal, wadding and silica gel sachet.

IDENTIFICATION OF SERENAGEN®

Brown, speckled, coated oval tablet.

REGISTRATION NUMBER –

To be allocated by SAHPRA upon registration.

ACCESS TO THE CORRESPONDING PROFESSIONAL INFORMATION

Scan QR code.

THIS LEAFLET WAS LAST REVISED ON

19 November 2024

NAME AND ADDRESS OF REGISTRATION HOLDER

Distributed by:

Amipro Advanced Development Products (Pty) Ltd
Unit 3, Eastgate Business Park, 1 South Road, Eastgate Extension,
Sandton, 2146
Tel: 011-802-8101



PASIËNT INLIGTINGSBLAD

SKEDULERINGSSTATUS ⁵⁰

Kategorie D: Komplementêre medisyne.

Dissipline: Dissipline Spesifiek

33.4 Tradisionele Chinese medisyne

HANDELSNAAM EN DOSEERVORM,

SERENAGEN[®], 60 en 120 tablette

LEES HIERDIE INLIGTINGSBLAD NOUKEURIG, WANT DIT BEVAT

BELANGRIKE INLIGTING VIR U

- **SERENAGEN[®]** is beskikbaar sonder doktersvoorskrif.
- U moet steeds **SERENAGEN[®]** versigtig gebruik om die beste resultate daaruit te kry.
- Hou hierdie inligtingsblad. Miskien moet u dit weer lees.
- Moenie **SERENAGEN[®]** met enige ander persoon deel nie.
- Vra jou apteker of gesondheidsorgpraktisyn of jy meer inligting of advies benodig.

WAT IS IN HIERDIE INLIGTINGSBLAD

- Wat **SERENAGEN[®]** is en waarvoor dit gebruik word.
- Wat jy moet weet voordat jy **SERENAGEN[®]** neem.
- Hoe om **SERENAGEN[®]** te neem.
- Moontlike nuwe-effekte.
- Hoe om **SERENAGEN[®]** te bêre.
- Inhoud van die pak en ander inligting

WATTER SERENAGEN[®] BEVAT

Elke twee (2) tablette **SERENAGEN[®]** bevat die volgende aktiewe:

Rehmannia (Rehmannia glutinosa) [Wortel 4:1 uittreksel]	290 mg
Schisandra (Schisandra chinensis) [Vrug 4:1 uittreksel]	120 mg
Jujube (Ziziphus spinosa) [Vrug 4:1 uittreksel]	120 mg
Chinese Asperge (Asparagus cochinchinensis) [Wortel 4:1 uittreksel]	70 mg
Dong Quai (Angelica sinensis) [Wortel 4:1 uittreksel]	70 mg
Ophiopogon (Ophiopogon japonicus) [Wortel 4:1 uittreksel]	70 mg
Scrophularia (Scrophularia ningpoensis) [Wortel 4:1 uittreksel]	70 mg
Asiatiese Ginseng (Panax ginseng) [Wortel 4:1 uittreksel]	40 mg
Chinese Salvia (Salvia miltiorrhiza) [Wortel 4:1 uittreksel]	40 mg
Poria Fungus (Wolfiporia cocos) [Hele 4:1 uittreksel]	40 mg
Polygala (Polygala tenuifolia) [Wortel 4:1 uittreksel]	40 mg
Platycodon (Platycodon grandiflorum) [Wortel 4:1 uittreksel]	30 mg

Onaktiewe bestanddele: Mikrokrystaliese sellulose, kroskarmellose-natrium, sellulose, steariensus (plant), silika, magnesiumstearaat (plant), en bedekking (hipromellose, mediumkietting triglisieriede, en hidroksipropielsellulose).

Hierdie produk is geskik vir vegetariërs, is nie-GGO, gluten en suiwelvry.

WAT IS SERENAGEN[®] KAPSULES EN WAARVOOR WORD DIT GEBRUIK
SERENAGEN[®] bied 'n unieke kombinasie van adaptogene kruie wat tradisioneel gebruik word om die organe en stelsels te ondersteun wat betrokke is by die liggaam se stresreaksie. Dit is geformuleer vir individue wat gestres en gespanne is.

VOORDAT JY SERENAGEN[®] NEEM

As jy enige chroniese medisyne neem, moenie hierdie produk gebruik sonder om jou gesondheidsorgpraktisyn te raadpleeg nie. Moenie **SERENAGEN[®]** neem as jy hipersensitief (allergies) is vir enige van die bestanddele nie. As jy allergieë het, maak seker dat jy met jou gesondheidsorgpraktisyn gaan voordat jy hierdie aanvulling neem.

Chirurgie: **SERENAGEN[®]** kan bloedsuikervlakke beïnvloed. Dit kan inname met bloedsuikerbeheer tydens en na die operasie. Hou op om **SERENAGEN[®]** ten minste 2 weke voor 'n geskeduleerde operasie te gebruik. **Latex allergie:** Mense wat allergies is vir latex kan ook allergies wees vir **SERENAGEN[®]**.

Hierdie medisyne is nie deur SAHPRA geëvalueer vir kwaliteit, veiligheid of beoogde gebruik nie.

SWANGERSKAP EN BORSVOEDING

Veiligheid tydens swangerskap en borsvoeding is nie vasgestel nie, raadpleeg u gesondheidsorgpraktisyn voor gebruik. As u swanger is of borsvoed, dink dat u swanger is of beplan om 'n baba te hê, raadpleeg u gesondheidsorgpraktisyn voordat u hierdie aanvulling neem.

BESTUUR EN GEBRUIK VAN MASHIENE

Dit is nie altyd moontlik om te voorspel in watter mate **SERENAGEN[®]** die daaglikse aktiwiteite van 'n pasiënt kan belemmer nie. Pasiënte moet seker maak dat hulle nie betrokke raak by aktiwiteite wat verband hou met die vereiste van verstandelike waaksaamheid, oordeel en/of gesonde koördinasie en visie nie, byvoorbeeld bestuur, vr, vlieg, seil, bedryfsmasjiene/toerusting, totdat hulle bewus is van die mate waartoe **SERENAGEN[®]** jou beïnvloed.

NEEM VAN ANDER MEDISYNE SAAM MET SERENAGEN[®]

Komplementêre medisyne produkte bevat bestanddele wat met sommige medisyne 'n interaksie kan hê. As jy enige chroniese medisyne gebruik, moenie hierdie produk gebruik sonder om jou gesondheidsorgpraktisyn te raadpleeg nie.

HOE OM SERENAGEN[®] TE NEEM

Moenie medisyne wat vir u voorgeskryf is, met enige ander persoon deel nie. Neem altyd **SERENAGEN[®]** presies soos beskryf in hierdie inligtingsblad, of soos u dokter, gesondheidsorgpraktisyn voorgeskryf het. Raadpleeg jou dokter, apteker of gesondheidsorgpraktisyn as jy onseker is.

Aanwysings vir gebruik:

Volwasse: Neem twee (2) tablette daaglik of soos voorgeskryf deur jou gesondheidspraktisyn.

Moenie meer as die aanbevole dosis neem nie.

AS JY MEER SERENAGEN[®] NEEM AS WAT JY MOET

Indien die geval van oordosering, raadpleeg u dokter of apteker. Indien nie een hiervan beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum.

AS JY VERGEET OM SERENAGEN[®] TE NEEM

Moenie 'n dubbele dosis neem om op te maak vir 'n dosis wat oorgeslaan is nie.

MOONTLIKE NUWE-EFFEKTE

SERENAGEN[®] kan nuwe-effekte hê.

Indien jy enige van die volgende gediagnoseerde toestande het, raadpleeg jou gesondheidsorgpraktisyn voor gebruik: Diabetes, hoë bloeddruk, bloedingsversteurings, hormoon sensitiewe toestande soos borskanker, baarmoederkanker, eierstokkanker, endometriose of baarmoederfibroïde. As jy die volgende medisyne neem, raadpleeg asseblief jou gesondheidspraktisyn voordat jy neem: diabetes (Antidiabetiese dwelms), hoë bloeddruk (Antihypertensiewe middels), Bloedingsversteurings, stadige bloedstolling (Antikoagulant / Antiplaatjie middels). As enige van die volgende gebeur, hou op om **SERENAGEN[®]** te gebruik en vertel jou dokter of gaan na die ongevalle-afdeling by jou naaste hospitaal:

- Allergiese reaksies-uitslag, jeuk / swelling (veral van die gesig / tong / keel), erge duiseligheid, probleme met asemhaling;

Dit is alles baie ernstige gevolge. As jy dit het, het jy dalk 'n ernstige reaksie op **SERENAGEN[®]** gehad. Jy mag dalk dringende mediese hulp of hospitalisasie nodig hê.

Nuwe-effekte wat gewoonlik nie mediese hulp benodig nie (rapporteur aan u gesondheidsorgpraktisyn as dit voortduur):

- naarheid / braking, hardlywigheid, verlies aan eetlust, verhoogde dors, verhoogde urinering, geestelike / gemoedsveranderinge, ongewone moegheid.

Nie alle nuwe-effekte en interaksies wat vir **SERENAGEN[®]** gerapporteur word, is by hierdie inligtingsblad gesluit nie. Indien jou algemene gesondheid vererger of as jy enige nadelige effekte ervaar tydens die gebruik van hierdie medisyne, raadpleeg asseblief jou dokter, apteker of ander gesondheidsorgpraktisyn vir raad. As jy enige nuwe-effekte sien wat nie in hierdie inligtingsblad genoem word nie, stel asseblief jou dokter, apteker of gesondheidsorgpraktisyn in kennis.

AANMELDING VAN NUWE-EFFEKTE

As jy nuwe-effekte ervaar, praat met jou dokter, apteker of ander gesondheidsorgpraktisyn. U kan ook nuwe-effekte aan SAHPRA rapporteer via "6.04 Adverse Drug Reaction Reporting Form", wat aanlyn gevind word onder SAHPRA se publikasies:

<https://www.sahpra.org.za/Publications/Index/8>. Deur nuwe-effekte aan te meld, kan u help om meer inligting te verskaf oor die veiligheid van **SERENAGEN[®]**.

HOE OM SERENAGEN[®] TE BÊRE

Bêre alle medisyne buite bereik van kinders.

Bêre in 'n koel, donker plek, onder 25 °C.

Beskermt teen direkte sonlig en hitte.

Bêre in die oorspronklike verpakking.

Moet nie gebruik word na die vervaldatum wat op die bottel aangedui word nie.

Moenie **SERENAGEN[®]** gebruik as u sigbare tekens van agteruitgang opmerk nie.

Moenie ongebruikte medisyne in dreine of rioolstelsels (bv. toilette) weggooi nie.

AANBIEDING VAN SERENAGEN[®]

60 TABLETTE - 'n Wit 175 ml HDPE-bottel met 'n wit dop, induksieseël, peuterkrimpseseël, spons en silikagelsakke.

120 TABLETTE - 'n Wit 250 ml HDPE-bottel met 'n wit dop, induksieseël, peuterkrimpseseël, spons en silikagelsakke.

IDENTIFIKASIE VAN SERENAGEN[®]
Bruin, gespikkelde, bedekte ovaal tablet.

REGISTRASIONOMMER

Om deur SAHPRA toegeken te word by registrasie.

TOEGANG TOT DIE OOREENSTEMMENDE PROFESIONELE

INLIGTING

Skandeer QR-kode..

HIERDIE INLIGTINGSBLAD IS LAAS HERSIEN OP

19 November 2024

NAAM EN ADRES VAN REGISTRASIEHOUER

Versprei deur:

Amipro Advanced Development Products (Pty) Ltd

Unit 3, Eastgate Business Park, 1 South Road, Eastgate Extension,

Sandton, 2146

Tel: 011-802-8101



