

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S0

Category D: Complementary medicine.

Discipline: Health Supplement

34.12 Multiple substance formulation

PROPRIETARY NAME, STRENGTH, PHARMACEUTICAL FORM –

TRANCOR[®], 60 & 120 Capsules

READ ALL OF THIS LEAFLET CAREFULLY BECAUSE IT CONTAINS IMPORTANT INFORMATION FOR YOU

- **TRANCOR[®]** is available without a doctor's prescription.
- Nevertheless, you still need to use **TRANCOR[®]** carefully to get the best results from it.
- Keep this leaflet. You may need to read it again.
- Do not share **TRANCOR[®]** with any other person.
- Ask your pharmacist or healthcare professional if you need more information or advice.

WHAT IS IN THIS LEAFLET

- What **TRANCOR[®]** is and what it is used for.
- What you need to know before you take **TRANCOR[®]**.
- How to take **TRANCOR[®]**.
- Possible side effects.
- How to store **TRANCOR[®]**.
- Contents of the pack and other information.

WHAT TRANCOR[®] CONTAINS

Each one (1) capsule of **TRANCOR[®]** contains the following actives:

Taurine	500 mg
L-Cysteine (as N-acetyl-L-cysteine)	300 mg
Magnesium (as Magnesium Malate)	100 mg
Decaffeinated Green Tea (Camellia sinensis) [Leaf]	150 mg
(Extract 25:1 ratio standardized to 60% (90 mg) catechins and 40% (60 mg) epigallocatechin gallate (EGCG))	
Vitamin B6 (as pyridoxine HCl)	12,5 mg

Inactive Ingredients: Capsule (hydroxypropylmethylcellulose), magnesium stearate (vegetable), microcrystalline cellulose and silica.

This product is suitable for Vegetarians and non-GMO. (Sugar Free)

WHAT TRANCOR[®] IS AND WHAT IT IS USED FOR

TRANCOR[®] is formulated to support a sense of calmness. This advanced formula features ingredients that cross the blood-brain barrier for targeted neurological support.

BEFORE YOU TAKE TRANCOR[®]

If you are taking any chronic medication, do not use this product without consulting your healthcare provider. Do not take **TRANCOR[®]** if you are hypersensitive (allergic) to any of the ingredients. If you have allergies, be sure to check with your healthcare professional before taking this supplement. **Surgery:** **TRANCOR[®]** might slow blood clotting. This might increase the risk of bleeding during and after surgery. Stop taking **TRANCOR[®]** at least 2 weeks before a scheduled surgery.

This medicine has not been evaluated by the SAHPRA for quality, safety, or intended use.

PREGNANCY AND BREASTFEEDING

Safety during pregnancy and breastfeeding has not been established, consult with your healthcare professional before use. If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare practitioner for advice before taking this supplement.

DRIVING AND USING MACHINES

It is not always possible to predict to what extent **TRANCOR[®]** may interfere with the daily activities of a patient. Patients should ensure that they do not engage in activities relating to requiring mental alertness, judgement and/or sound co-ordination and vision e.g. driving, riding, flying, sailing, operating machines/equipment, until they are aware of the measure to which **TRANCOR[®]** may affect you.

TAKING OTHER MEDICINES WITH TRANCOR[®]

Complementary medicine products contain ingredients which may interact with some medication. If you are taking any chronic medication, do not use this product without consulting your medical practitioner

HOW TO TAKE TRANCOR[®]

Do not share medicines prescribed for you with any other person. Always take **TRANCOR[®]** exactly as described in this leaflet, or as your doctor, healthcare provider has instructed you. You should check with your doctor, pharmacist or healthcare provider if you are unsure.

Directions for use:

Adults: Take two (2) capsules twice daily or as directed by your healthcare practitioner.

Do not take more than the recommended dose.

IF YOU TAKE MORE TRANCOR[®] THAN YOU SHOULD

In the event of overdose, consult your doctor, pharmacist. Should none of these be available, contact the nearest hospital or poison control centre.



IF YOU FORGET TO TAKE TRANCOR[®]

Do not take a double dose to make up for a missed dose.

POSSIBLE SIDE EFFECTS

TRANCOR[®] may have side effects.

Should you have any of the following diagnosed conditions, consult with your healthcare professional before use: Asthma, Bleeding disorders, Anemia, Heart conditions, Diabetes, Glaucoma, High blood pressure, Irritable bowel syndrome (IBS), Liver disease, Weak bones (osteoporosis), Post-surgical stent placement, Weight loss surgery. If you are taking the following medication, please consult with your healthcare practitioner before taking: Lithium; Medications for high blood pressure (Antihypertensive drugs), Nitroglycerin, Activated charcoal, Chloroquine (Aralen), Medications for high blood pressure (Antihypertensive drugs), Medications that slow blood clotting (Anticoagulant / Antiplatelet drugs), Ephedrine, Atorvastatin (Lipitor), Medications that can harm the liver (Hepatotoxic drugs), Warfarin (Coumadin), Phenytoin (Dilantin), Levodopa. If any of the following happens, stop using **TRANCOR[®]** and tell your doctor or go to the casualty department at your nearest hospital:

- Allergic reactions – rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

These are all very serious effects. If you have them, you may have had a serious reaction to **TRANCOR[®]**. You may need urgent medical attention or hospitalisation.

Side effects that usually do not require medical attention (report to your healthcare professional if they continue):

- nausea, stomach pain, loss of appetite.

Not all side effects and interactions reported for **TRANCOR[®]** are included in this leaflet.

Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your doctor, pharmacist or other healthcare professional for advice. If you notice any side effects not mentioned in this leaflet, please inform your doctor, pharmacist or healthcare professional.

REPORTING OF SIDE EFFECTS

If you experience side effects, talk to your doctor, pharmacist or other healthcare professional. You can also report side effects to SAHPRA via "6.04 Adverse Drug Reaction Reporting Form", found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of **TRANCOR[®]**.

STORAGE AND DISPOSING OF TRANCOR[®]

Store all medicines out of the reach of children.

Store in a cool, dark place, below 25 °C.

Protect from direct sunlight and heat.

Store in the original packaging.

Do not use after the expiry date stated on the bottle.

Do not use **TRANCOR[®]** if you notice visible signs of deterioration.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

PRESENTATION OF TRANCOR[®]

60 CAPSULES - A white 175 ml HDPE bottle with a white cap, induction seal, tamper shrink seal, wadding and silica gel sachet.

120 CAPSULES - A white 250 ml HDPE bottle with a white cap, induction seal, tamper shrink seal, wadding and silica gel sachet.

IDENTIFICATION OF TRANCOR[®]

Clear vegetable capsule with tan speckled powder.

REGISTRATION NUMBER –

To be allocated by SAHPRA upon registration.

ACCESS TO THE CORRESPONDING PROFESSIONAL INFORMATION

Scan QR code.

THIS LEAFLET WAS LAST REVISED ON

19 November 2024

NAME AND ADDRESS OF REGISTRATION HOLDER

Distributed by:

Amipro Advanced Development Products (Pty) Ltd
Unit 3, Eastgate Business Park, 1 South Road, Eastgate Extension,
Sandton, 2146
Tel: 011-802-8101



PASIËNT INLIGTINGSBLAD

SKEDULERINGSSTATUS **50**

Kategorie D: Komplementêre medisyne.

Dissipline: Gesondheidsaanvulling

34.12 Meervoudige Bestandeel Formulasie

HANDELSNAAM EN DOSEERVORM,

TRANCOR[®], 60 & 120 Kapsule

LEES HIERDIE INLIGTINGSBLAD NOUKEURIG, WANT DIT BEVAT

BELANGRIKE INLIGTING VIR U

- **TRANCOR[®]** is beskikbaar sonder doktersvoorskrif.
- U moet steeds **TRANCOR[®]** versigtig gebruik om die beste resultate daaruit te kry.
- Hou hierdie inligtingsblad. Miskien moet u dit weer lees.
- Moenie **TRANCOR[®]** met enige ander persoon deel nie.
- Vra jou apteker of gesondheidsorgpraktisyn of jy meer inligting of advies benodig.

WAT IS IN HIERDIE INLIGTINGSBLAD

- Wat **TRANCOR[®]** is en waarvoor dit gebruik word.
- Wat jy moet weet voordat jy **TRANCOR[®]** neem.
- Hoe om **TRANCOR[®]** te neem.
- Moontlike nuwe-effekte.
- Hoe om **TRANCOR[®]** te bêre.
- Inhoud van die pak en ander inligting

WAT TRANCOR[®] BEVAT

Elke een (1) kapsule **TRANCOR[®]** bevat die volgende aktiewe:

Taurine	500 mg
L-sisteien (as N-asetiel-L-sisteien)	300 mg
Magnesium (as magnesiummalaat)	100 mg
Kafeienvrye groen tee (Camellia sinensis) [blaar]	150 mg
(Uittreksel 25: 1 verhouding gestandaardiseer tot 60% (90 mg) katekiene en 40%) (60 mg) epigallocatechin gallate (EGCG))	
Vitamine B6 (as piridoksien HCl)	12,5 mg

Onaktiewe bestanddele Kapsule (hidroksiopropylmetiellselulose), magnesiumstearaat (groente), mikrokristallyne selulose en silika.

Hierdie produk is geskik vir vegetariërs en nie-GGO. (Suikervry)

WAT IS TRANCOR[®] EN WAARVOOR WORD DIT GEBRUIK

TRANCOR[®] is geformuleer om 'n gevoel van kalmte te ondersteun. Hierdie gevorderde formule bevat bestanddele wat die bloedreïnsperping oorsake vir geteikende neurologiese ondersteuning.

VOORDAT JY TRANCOR[®] NEEM

As jy enige chroniese medisyne neem, moenie hierdie produk gebruik sonder om jou gesondheidsorgpraktisyn te raadpleeg nie. Moenie **TRANCOR[®]** neem as jy hipersensitief (allergies) is vir enige van die bestanddele nie. As jy allergieë het, maak seker dat jy met jou gesondheidsorgpraktisyn gaan voordat jy hierdie aanvulling neem. **Chirurgie:** **TRANCOR[®]** kan bloedstolling vertraag. Dit kan die risiko van bloeding tydens en na die operasie verhoog. Hou op om **TRANCOR[®]** ten minste 2 weke voor 'n geskeduleerde operasie te neem.

Hierdie medisyne is nie deur SAHPRA geëvalueer vir kwaliteit, veiligheid of beoogde gebruik nie.

SWANGERSKAP EN BORSVOEDING

Veiligheid tydens swangerskap en borsvoeding is nie vasgestel nie, raadpleeg u gesondheidswerker voor gebruik. As u swanger is of borsvoed, dink dat u swanger is of beplan om 'n baba te hê, raadpleeg u gesondheidsorgpraktisyn voordat u hierdie aanvulling neem.

BESTUUR EN GEBRUIK VAN MASIËNE

Dit is nie altyd moontlik om te voorspel in watter mate **TRANCOR[®]** die daaglikse aktiwiteite van 'n pasiënt kan belemmer nie. Pasiënte moet seker maak dat hulle nie betrokke raak by aktiwiteite wat verband hou met die vereiste van verstandelike waaksaamheid, oordeel en/of gesonde koördinasie en visie nie, byvoorbeeld bestuur, ry, vlieg, seil, bedryfsmasjiene/toerusting, totdat hulle bewus is van die mate waartoe **TRANCOR[®]** jou beïnvloed.

NEEM VAN ANDER MEDISYNE SAAM MET TRANCOR[®]

Komplementêre medisyne produkte bevat bestanddele wat met sommige medisyne 'n interaksie kan hê. As jy enige chroniese medisyne gebruik, moenie hierdie produk gebruik sonder om jou gesondheidsorgpraktisyn te raadpleeg nie.

HOE OM TRANCOR[®] TE NEEM

Moenie medisyne wat vir u voorgeskryf is, met enige ander persoon deel nie. Neem altyd **TRANCOR[®]** presies soos beskryf in hierdie inligtingsblad, of soos u dokter, gesondheidsorgpraktisyn voorgeskryf het. Raadpleeg jou dokter, apteker of gesondheidsorgpraktisyn as jy onseker is.

Aanwysings vir gebruik:

Volwassenes: Neem twee (2) kapsules twee keer per dag of soos voorgeskryf deur jou gesondheidsorgpraktisyn.

Moenie meer as die aanbevole dosis neem nie.

AS JY MEER TRANCOR[®] NEEM AS WAT JY MOET

In die geval van oordosering, raadpleeg u dokter of apteker. Indien nie een hiervan beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum.

AS JY VERGEET OM TRANCOR[®] TE NEEM

Moenie 'n dubbele dosis neem om te maak vir 'n dosis wat oorgeslaan is nie.

MOONTLIKE NUWE-EFFEKTE

TRANCOR[®] kan nuwe-effekte hê.

Indien jy enige van die volgende gediagnoseerde toestande het, raadpleeg jou gesondheidsorgpraktisyn voor gebruik: Asma, bloedingsversteurings, bloedarmoede, harttoestande, diabetes, gloukoom, hoë bloeddruk, prikkelbare dermsindroom (IBS), lewersiekte, swak bene (osteoporose), post-chirurgiese stentplasing, gewigsverlieschirurgie. As jy die volgende medisyne neem, raadpleeg asseblief jou gesondheidspraktisyn voordat jy neem: Litium, medisyne vir hoë bloeddruk (Antihipertensiewe middels), Nitroglyserin, Geaktiveerde houtskool, Chlorokien (Aralen), medisyne vir hoë bloeddruk (Antihipertensiewe middels), medisyne wat bloedstolling vertraag (Antikoagulanter / Antiplatelet-middels), Efedrien, Atorvastatin (Lipitor), medisyne wat die lewer kan benadeel (Hepatotoksiese middels), Warfarin (Coumadin), Fenitioien (Dilantin), Levodopa. As enige van die volgende gebeur, hou op om **TRANCOR[®]** te gebruik en vertel jou dokter of gaan na die ongevalle-afdeling by jou naaste hospitaal:

- Allergiese reaksies: uitslag, jeuk / swelling (veral van die gesig / tong / keel), erge duiseligheid, probleme met asemhaling;

Dit is alles baie ernstige gevolge. As jy dit het, het jy dalk 'n ernstige reaksie op **TRANCOR[®]** gehad. Jy mag dalk dringende mediese hulp of hospitalisasie nodig hê.

Nuwe-effekte wat gewoonlik nie mediese hulp benodig nie (rapporteur aan u gesondheidsorgpraktisyn as dit voortduur):

- naarheid, maagpyn, verlies aan eetlust.

Nie alle nuwe-effekte en interaksies wat vir **TRANCOR[®]** gerapporteer word, is by hierdie inligtingsblad ingesluit nie. Indien jou algemene gesondheid vererger of as jy enige nadelige effekte ervaar tydens die gebruik van hierdie medisyne, raadpleeg asseblief jou dokter, apteker of ander gesondheidsorgpraktisyn vir raad. As jy enige nuwe-effekte sien wat nie in hierdie inligtingsblad genoem word nie, stel asseblief jou dokter, apteker of gesondheidsorgpraktisyn in kennis.

AANMELDING VAN NUWE-EFFEKTE

As jy nuwe-effekte ervaar, praat met jou dokter, apteker of ander gesondheidsorgpraktisyn. U kan ook nuwe-effekte aan SAHPRA rapporteer via "6.04 Adverse Drug Reaction Reporting Form", wat aanlyn gevind word onder SAHPRA se publikasies:

<https://www.sahpra.org.za/Publications/Index/8>. Deur nuwe-effekte aan te meld, kan u help om meer inligting te verskaf oor die veiligheid van **TRANCOR[®]**.

HOE OM TRANCOR[®] TE BÊRE

Bêre alle medisyne buite bereik van kinders.

Bêre in 'n koel, donker plek, onder 25 °C.

Beskermt teen direkte sonlig en hitte.

Bêre in die oorspronklike verpakking.

Moet nie gebruik word na die vervaldatum wat op die bottel aangedui word nie.

Moenie **TRANCOR[®]** gebruik as u sigbare tekens van agteruitgang opmerk nie.

Moenie ongebruikte medisyne in dreine of rioolstelsels (bv. toilette) weggooi nie.

AANBIEDING VAN TRANCOR[®]

60 KAPSULES - 'n Wit 175 ml HDPE-bottel met 'n wit dop, induksiesêl, peuterkrimpsêl, spons en silikagelsakke.

120 TABLETTE - 'n Wit 250 ml HDPE-bottel met 'n wit dop, induksiesêl, peuterkrimpsêl, spons en silikagelsakke.

IDENTIFIKASIE VAN TRANCOR[®]

Duidelik groentekapsule met bruin gespikkelde poeier.

REGISTRASIONUMMER

Om deur SAHPRA toegeken te word by registrasie.

TOEGANG TOT DIE OOREENSTEMMENDE PROFESSIONELE INLIGTING

Skandeer QR-kode.

HIERDIE INLIGTINGSBLAD IS LAAS HERSIEN OP

19 November 2024

NAAM EN ADRES VAN REGISTRASIEHOUER

Versprei deur:

Amipro Advanced Development Products (Pty) Ltd

Unit 3, Eastgate Business Park, 1 South Road, Eastgate Extension,

Sandton, 2146

Tel: 011-802-8101

